

Durham Research Online

Deposited in DRO:

12 November 2014

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Magni, V. and Faccenna, C. and van Hunen, J. and Funicello, F. (2014) 'How collision triggers backarc extension : insight into Mediterranean style of extension from 3-D numerical models.', *Geology*, 42 (6). pp. 511-514.

Further information on publisher's website:

<http://dx.doi.org/10.1130/G35446.1>

Publisher's copyright statement:

© 2014 Geological Society of America. Gold Open Access: This paper is published under the terms of the CC-BY license.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

How collision triggers backarc extension: Insight into Mediterranean style of extension from 3-D numerical models

Valentina Magni¹, Claudio Faccenna², Jeroen van Hunen¹, and Francesca Funiciello²

¹Department of Earth Sciences, University of Durham, Durham DH1 3LE, UK

²Dipartimento di Scienze, Università Roma Tre, Largo San Leonardo Murialdo 1, 00146 Rome, Italy

ABSTRACT

The formation and evolution of a backarc basin are linked to the dynamics of the subduction system. The opening of the central Mediterranean basins is a well-documented example of backarc extension characterized by short-lived episodes of spreading. The underlying reasons for this episodicity are obscured by the complexity of this subduction system, in which multiple continental blocks enter the subduction zone. We present results from three-dimensional numerical models of laterally varying subduction to explain the mechanism of backarc basin opening and the episodic style of spreading. Our results show that efficient backarc extension can be obtained with an along-trench variation in slab buoyancy that produces localized deformation within the overriding plate. We observe peaks in the trench retreating velocity corresponding first to the opening of the backarc basin, and later to the formation of slab windows. We suggest that the observed episodic trench retreat behavior in the central Mediterranean is caused by the formation of slab windows.

INTRODUCTION

Backarc basins (BABs) form in convergent margins where the overriding plate is in an extensional regime (e.g., Uyeda and Kanamori, 1979; Carlson and Melia, 1984). Their formation is kinematically associated either with trench retreat (e.g., Japan, Lau-Kermadec, and Philippine basins) or with motion of the upper plate away from the trench (e.g., the Mariana Islands). What triggers backarc extension and why it is episodic (lasting only for tens of millions of years; e.g., Karig, 1971) are debated. In several cases, such as the Woodlark Basin (Papua New Guinea) or Havre Trough BAB (New Zealand), it has been noted that backarc extension takes place by an adjacent collisional zone inducing sharp rotation and curvature of the adjacent region (Wallace et al., 2009). Such a setting is typical for the Mediterranean. In the eastern Mediterranean, for example, backarc extension in the Aegean mainly occurred during Arabia-Eurasia collision and accelerated after the associated slab tearing along western Turkey in the east, and near Corinth in the west (Faccenna et al., 2006; Jolivet et al., 2013). Moreover, in the central Mediterranean, the retreat of the Calabrian slab led to the formation of two main BABs. Geological records show how this process occurred episodically, with pulses of greater extension and oceanic spreading (e.g., Malinverno and Ryan, 1986; Faccenna et al., 2001a) (Fig. 1). The first episode of extension occurred between 30 and 16 Ma and led to the formation of the Liguro-Provençal basin. After a pause of ~5 m.y., extension shifted to the east of Sardinia to form the Tyrrhenian basin (where rifting started ca. 12–10 Ma). Here, pulses of oceanic spreading migrated eastward and formed the Vavilov Basin (ca. 4.3–2.6 Ma; Kastens and Mascle, 1990), followed by the Marsili

Basin (ca. 2 Ma; Patacca et al., 1990; Nicolosi et al., 2006). At depth, the slab broke off and tore where collision occurred; first in northern Africa, ca. 10–12 Ma, as evidenced by the change in the nature of volcanism of the Mogods volcanic complex (Faccenna et al., 2005), and later in the Apennines. The lateral tearing of the slab along the continental margins adjacent to the oceanic subduction accommodates the Calabrian slab rollback (Carminati et al., 1998; Wortel and Spakman, 2000). The present-day configuration of the central Mediterranean subduction zone is supported by tomographic images (Piomallo and Morelli, 2003; Fig. 1) that show two slab windows next to the narrow Calabrian slab. The nature of the episodic behavior of BAB opening is debated. It has been postulated that slowing of extension can be driven by buoyant continental crust entering the trench (Malinverno and Ryan, 1986), by the interaction between the slab tip and the 660 km seismic discontinuity (Faccenna et al., 2001b), or by the collision-related lateral slab tear and subsequent development of vigorous toroidal flow around the slab (Faccenna et al., 2005, 2007; Guillaume et al., 2010).

Inspired by the geological cases described here, and in particular by the Mediterranean example, we developed three-dimensional (3-D) numerical models to explore how laterally varying collision may influence the formation and style of backarc extension. Our results show that BABs form next to collisional systems and that they develop episodically if the boundary between oceanic and continental lithosphere ruptures.

MODELS OF CONTINENTAL COLLISION AND BACKARC BASIN DEVELOPMENT

Previous studies helped our understanding of the mechanisms of the BAB formation through

2-D (Arcay et al., 2008; Gerya and Meilick, 2011) and laterally heterogeneous 3-D models (Capitanio and Replumaz, 2013; Li et al., 2013), and the episodicity in backarc tectonic regimes (Clark et al., 2008; Baitsch-Ghirardello et al., 2014). However, until now, all these aspects have been studied separately.

We present a set of 3-D numerical models of subduction using the finite element code Citcom (Moresi and Gurnis, 1996; Zhong et al., 2000; van Hunen et al., 2005) to explore the role of collision on BAB opening. Conservation of mass, momentum, thermal energy, and composition describe material flow, deformation, and thermal state. Continental collision is simulated using a viscoplastic rheology that combines diffusion creep, dislocation creep, and a pseudo-brittle deformation simulating Byerlee's law (van Hunen and Allen, 2011), while the converging plates are decoupled through a mobile weak zone (Magni et al., 2012; see the GSA Data Repository¹ for details on the numerical method).

To investigate the role of lateral continental colliding blocks on the side of an oceanic setting, we vary the width of the subducting oceanic plate from 500 to 1200 km (Fig. 2) within a 3300 × 3960 × 660 km regional 3-D model, in which the overriding plate is fully continental, and the subducting plate is oceanic in the middle and continental at the sides (Fig. 2). For every model we compute the potential energy release rate (E'_{pot}) and the energy dissipation rate (E'_{diss}) to investigate how energy is distributed within the system (for details on how energy is computed, see the Data Repository). The models are not designed to simulate specifically any particular case, but rather to capture the dynamics of the system. All of these models show a common behavior during the first phase of evolution, when only oceanic lithosphere subducts and the entire slab slowly rolls back, causing trench retreat. When the continental blocks arrive at the trench, subduction locally slows and the trench starts to advance, while the oceanic part of the plate continues to subduct and sustains trench retreat. These opposite trench motions cause large strains in the overriding plate

¹GSA Data Repository item 2014181, details on the numerical method, energy computation, and distribution, is available online at www.geosociety.org/pubs/ft2014.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.




Figure 1. A: Reconstruction of central Mediterranean subduction system and trench retreating velocity computed from the amount of extension estimated in time along cross section shown in gray (in panel labeled “present day”; Faccenna et al., 2001a). Negative trench velocity indicates trench retreat. B: Isosurface of +0.8% P-wave velocity anomalies beneath central Mediterranean from tomography data (Piromallo and Morelli, 2003).

and a strong curvature of the trench and slab. The largest values of E'_{diss} in the overriding plate are mainly localized where ongoing oceanic subduction causes significant overriding plate deformation (Fig. 3A). For oceanic plates larger than 500 km, the overriding plate behind the oceanic subduction stretches and thins to accommodate the retreat until a BAB is formed ~10 m.y. after initial collision (Fig. 3B). During this process, the retreating velocity increases rapidly, from 1–2 cm/yr to 4–5.5 cm/yr in just a few million years (Fig. 4A). After the opening of the BAB, E'_{pot} integrated over the entire lithosphere slightly increases in response to this episode of faster trench retreat and associated slab sinking (Fig. 4B). This is because the sinking of cold, dense slabs is responsible for the

bulk of the potential energy release that powers the subduction dynamics (and, more generally, plate tectonics). Meanwhile, the E'_{diss} in the overriding plate drops (Fig. 4B), because most energy is now dissipated in the rapidly deforming subducting plate. At depth, the slab is highly deformed with a curved concave shape. The middle, oceanic part of the plate is still subducting, and slab pull is still acting there, whereas near the continental blocks, subduction almost stops and the slab starts breaking off in a necking process. At this point, rollback slows. If the oceanic plate is wider than 800 km, the oceanic slab pull is large enough to stretch and rupture the slab at the ocean-continent boundaries at a depth of ~230 km (Fig. 3D). These ruptures then rapidly propagate laterally, allowing, in only a




Figure 2. Initial model setup illustrating geometry, mechanical and thermal boundary conditions, and different lithologies. $T_m = 1350^\circ\text{C}$ is the mantle reference temperature and $\eta_m = 10^{20}\text{ Pa}\cdot\text{s}$ is the mantle reference viscosity (see the Data Repository [see footnote 1]).

few million years, the formation of two wide slab windows, where slab mantle material can flow through (Fig. 3E). This process corresponds to a second peak of acceleration in the trench retreat velocity (Fig. 4A) and a peak of increased E'_{pot} (Fig. 4B) that are not observed in narrower oceanic plates. The remaining oceanic slab becomes narrower (~400 km wide in the final stages) and the retreat slows.

A very different behavior is observed for the model with a 500-km-wide or narrower oceanic plate, where the retreat is very slow (<1 cm/yr), the slab deforms slightly, and no BAB forms.

We put the behavior of subducting ocean-continent plates in a wider context by investigating simpler models without along-strike variation in the subducting plate. Subduction of a purely oceanic plate (i.e., no continents at the sides) results in a very slow trench retreat and no BAB formation (Fig. DR1 in the Data Repository). In a model with one large continental block colliding along the entire trench after the subduction of oceanic lithosphere, the initial phase is very similar to the previous models. However, once collision occurs, subduction slows and the slab starts to advance slowly (Fig. 4A). At this point, the break-off starts in the center of the slab ~19 m.y. after collision and quickly propagates laterally (van Hunen and Allen, 2011) until the entire slab detaches from the shallow part of the plate. In both models without lateral variation (purely oceanic and laterally uniform continental lithosphere), no BAB is formed.

DISCUSSION AND CONCLUSIONS

These models highlight a number of essential ingredients for the formation of BABs. Localized deformation arising from rheological heterogeneities seems to be a key factor in the formation of a BAB. The model with only oceanic subduction shows that although the viscous dissipation in the overriding plate is high, it is not localized enough to open a BAB. In such a setting, BAB formation is still possible through other physical factors that are not taken into account in our models, such as the





Figure 3. Evolution of model with 1000-km-wide oceanic plate. A–C: Opening of the backarc basin, including plan views of distribution of energy dissipation rate (E'_{diss}) at surface and two x - z sections of the temperature field of continental subduction (a) and oceanic subduction (b). D, E: Slab tearing and slab window formation, including three-dimensional view of the slab (isosurface of $T = 1080$ °C) and the velocity field of the mantle in x - y section at middle of box ($z = 330$ km).



weakening of the overriding plate due to the presence of fluids and melt (Arcay et al., 2008; Gerya and Meilick, 2011).

When a continental block collides, the trench motion changes from retreating to advancing with respect to the upper plate and, where the trench remains parallel to the margin, the slab keeps advancing for few million years after collision (Magni et al., 2012). In the case of a continent of a finite size, a torque will be created between the continental advancing sides and the oceanic retreating one. The resulting localized tensile stresses in the overriding plate provide a means for localized stretching, which can lead to the opening of a BAB. The larger the oceanic plate, the greater the potential energy available to the system (Fig. 4B). Therefore, larger oceanic plates produce a larger

and faster rollback, rapidly inducing a BAB, whereas smaller oceanic plates may result in only a small amount of retreat, which is not vigorous enough to produce a BAB.

Slab pull also plays a crucial role in the formation of the slab windows at the ocean-continent boundaries, which are observed in the models only for a wide oceanic plate. Similar to BAB formation, slab tears at ocean-continent boundaries form under high tensile stresses, and can be generated only if the amount of slab pull is high enough. Once the oceanic slab has started tearing away from the continental block, it reinvigorates its retreating course, leading to a second trench retreat phase.

The numerical model setup is, by definition, a simplified and more generic version of reality. As such, the shape and geometry of the continental block are not designed to specifically reproduce the natural system as in the central Mediterranean. However, these models may serve to help us understand the dynamics of this complex subduction zone. There, the first phase of backarc extension ends when continental lithosphere enters the trench on both sides of the oceanic slab while the width of the oceanic domain narrows (Fig. 1). The onset of the second phase of extension occurs only a few million years after the first pulse. The first slab window forms ca. 12–10 Ma in northern Africa, and propagates laterally westward beneath Sicily until the middle Pleistocene, opening a 500-km-

Figure 4. A: Trench velocity in a y -section in the middle of the computational model domain (negative velocity: trench moves toward subducting plate). B: Potential energy (E'_{pot}) of lithosphere integrated over entire model domain. C: Energy dissipation (E'_{diss}) of overriding plate integrated over entire model domain of all the models (on right: only oc—model with purely oceanic subduction; oc + cont—model with oceanic subduction followed by laterally uniform continental lithosphere; o500—model with oceanic plate width of 500 km).

wide slab window. In the middle Pleistocene, a subduction window forms beneath the Central Apennines. Our model indicates that the opening of those slab windows is a necessary condition for the second phase of extension in the Tyrrhenian that lasted for ~5–7 m.y. The opening of the central Mediterranean is somewhat different from the simplified, perfectly symmetric numerical model presented here, but the rates, time scales, and dimensions of the model dynamics are quite similar to the natural example, indicating that the model represents the primary features of subduction forces, trench retreat, and slab deformation and tearing in the central Mediterranean. We furthermore propose that the presented model can, more generally, fit several other natural cases of backarc spreading associated with lateral variations along trench (such as the eastern Mediterranean and western Pacific Ocean). As in the case presented here, in these regions fast backarc extension is accompanied by large-scale rotation, caused by lateral collision of positively buoyant indentors, such as continental plates or plateaus (e.g., Tonga, Aegean, Papua New Guinea, New Zealand, the Mariana Islands, Vanuatu, and Fiji), ridges (e.g., the Ryukyu Islands), and seamount chains (e.g., Tonga) (Wallace et al., 2009, and references therein).

ACKNOWLEDGMENTS

We thank the editor J.B. Murphy and reviewers T. Gerya, W. Spakman, and S. Ellis for their constructive comments. This research was supported by the European Young Investigators (EURYI) Awards Scheme (EUROHORCS/ESF, including funds from the National Research Council of Italy) and by the European Research Council (ERC Starting Grant 279828). Models presented in this paper have been realized thanks to CASPUR HPC Standard Grant 2012.

REFERENCES CITED

- Arcay, D., Lallemand, S., and Doin, M.P., 2008, Back-arc strain in subduction zones: Statistical observations versus numerical modeling: *Geochemistry Geophysics Geosystems*, v. 9, Q02007, doi:10.1029/2007GC001875.
- Baitech-Ghirardello, B., Gerya, T.V., and Burg, J.-P., 2014, Geodynamic regimes of intra-oceanic subduction: Implications for arc extension vs. shortening processes: *Gondwana Research*, v. 25, p. 546–560, doi:10.1016/j.gr.2012.11.003.
- Capitanio, F.A., and Replumaz, A., 2013, Subduction and slab breakoff controls on Asian indentation tectonics and Himalayan western syntaxis formation: *Geochemistry Geophysics Geosystems*, v. 14, p. 3515–3531, doi:10.1002/ggge.20171.
- Carlson, R.L., and Melia, P.J., 1984, Subduction hinge migration: *Tectonophysics*, v. 102, p. 399–411, doi:10.1016/0040-1951(84)90024-6.
- Carminati, E., Wortel, M.J.R., Spakman, W., and Sabadini, R., 1998, The role of slab detachment processes in the opening of the western-central Mediterranean basins: Some geological and geophysical evidence: *Earth and Planetary Science Letters*, v. 160, p. 651–665, doi:10.1016/S0012-821X(98)00118-6.
- Clark, S.R., Stegman, D., and Müller, R.D., 2008, Episodicity in back-arc tectonic regimes: Physics of the Earth and Planetary Interiors, v. 171, p. 265–279, doi:10.1016/j.pepi.2008.04.012.
- Faccenna, C., Becker, T.W., Lucette, F.P., Jolivet, L., and Rossetti, F., 2001a, History of subduction and back-arc extension in the central Mediterranean: *Geophysical Journal International*, v. 145, p. 809–820, doi:10.1046/j.0956-540x.2001.01435.x.
- Faccenna, C., Funicello, F., Giardini, D., and Lucette, P., 2001b, Episodic backarc extension during restricted mantle convection in the central Mediterranean: *Earth and Planetary Science Letters*, v. 187, p. 105–116, doi:10.1016/S0012-821X(01)00280-1.
- Faccenna, C., Civetta, L., D'Antonio, M., Funicello, F., Margheriti, L., and Piromallo, C., 2005, Constraints on mantle circulation around the deforming Calabrian slab: *Geophysical Research Letters*, v. 32, L06311, doi:10.1029/2004GL021874.
- Faccenna, C., Bellier, O., Martinod, J., Piromallo, C., and Regard, V., 2006, Slab detachment beneath eastern Anatolia: A possible cause for the formation of the North Anatolian fault: *Earth and Planetary Science Letters*, v. 242, p. 85–97, doi:10.1016/j.epsl.2005.11.046.
- Faccenna, C., Funicello, F., Civetta, L., D'Antonio, M., Moroni, M., and Piromallo, C., 2007, Slab disruption, mantle circulation, and the opening of the Tyrrhenian basins, in Beccaluva, L., ed., *Cenozoic volcanism in the Mediterranean area: Geological Society of America Special Paper* 418, p. 153–169, doi:10.1130/2007.2418(08).
- Gerya, T.V., and Meilick, F.I., 2011, Geodynamic regimes of subduction under an active margin: Effects of rheological weakening by fluids and melts: *Journal of Metamorphic Geology*, v. 29, p. 7–31, doi:10.1111/j.1525-1314.2010.00904.x.
- Guillaume, B., Funicello, F., Faccenna, C., Martinod, J., and Olivetti, V., 2010, Spreading pulses of the Tyrrhenian Sea during the narrowing of the Calabrian slab: *Geology*, v. 38, p. 819–822, doi:10.1130/G31038.1.
- Jolivet, L., and 22 others, 2013, Aegean tectonics: Strain localisation, slab tearing and trench retreat: *Tectonophysics*, v. 597–598, p. 1–33, doi:10.1016/j.tecto.2012.06.011.
- Karig, D.E., 1971, Origin and development of marginal basins in the western Pacific: *Journal of Geophysical Research*, v. 76, p. 2542–2561, doi:10.1029/JB076i011p02542.
- Kastens, K.A., and Mascle, J., 1990, The geological evolution of the Tyrrhenian Sea: An introduction to the scientific results of ODP Leg 107, in Kastens, K.A., et al., *Proceedings of the Ocean Drilling Program, Scientific results, Volume 107: College Station, Texas, Ocean Drilling Program*, p. 3–26, doi:10.2973/odp.proc.sr.107.187.1990.
- Li, Z.-H., Xu, Z., Gerya, T., and Burg, J.-P., 2013, Collision of continental corner from 3-D numerical modeling: *Earth and Planetary Science Letters*, v. 380, p. 98–111, doi:10.1016/j.epsl.2013.08.034.
- Magni, V., van Hunen, J., Funicello, F., and Faccenna, C., 2012, Numerical models of slab migration in continental collision zones: *Solid Earth*, v. 3, p. 293–306, doi:10.5194/se-3-293-2012.
- Malinverno, A., and Ryan, W.B.F., 1986, Extension in the Tyrrhenian Sea and shortening in the Apennines as result of arc migration driven by sinking of the lithosphere: *Tectonics*, v. 5, p. 227–245, doi:10.1029/TC005i002p00227.
- Moresi, L., and Gurnis, M., 1996, Constraints on the lateral strength of slabs from three-dimensional dynamic flow models: *Earth and Planetary Science Letters*, v. 138, p. 15–28, doi:10.1016/0012-821X(95)00221-W.
- Nicolosi, I., Speranza, F., and Chiappini, M., 2006, Ultrafast oceanic spreading of the Marsili Basin, southern Tyrrhenian Sea: Evidence from magnetic anomaly analysis: *Geology*, v. 34, p. 717–720, doi:10.1130/G22555.1.
- Patacca, E., Sartori, R., and Scandone, P., 1990, Tyrrhenian Basin and Apenninic arcs: Kinematic relations since late Tortonian times: *Memorie della Società Geologica Italiana*, v. 45, p. 425–451.
- Piromallo, C., and Morelli, A., 2003, P wave tomography of the mantle under the Alpine-Mediterranean area: *Journal of Geophysical Research*, v. 108, no. B2, p. 1–22, doi:10.1029/2002JB001757.
- Uyeda, S., and Kanamori, H., 1979, Back-arc opening and the mode of subduction: *Journal of Geophysical Research*, v. 84, p. 1049–1061, doi:10.1029/JB084iB03p01049.
- van Hunen, J., and Allen, M.B., 2011, Continental collision and slab break-off: A comparison of 3-D numerical models with observations: *Earth and Planetary Science Letters*, v. 302, p. 27–37, doi:10.1016/j.epsl.2010.11.035.
- van Hunen, J., Zhong, S., Shapiro, N.M., and Ritzwoller, M.H., 2005, New evidence for dislocation creep from 3-D geodynamic modeling of the Pacific upper mantle structure: *Earth and Planetary Science Letters*, v. 238, p. 146–155, doi:10.1016/j.epsl.2005.07.006.
- Wallace, L.M., Ellis, S., and Mann, P., 2009, Collisional model for rapid fore-arc block rotations, arc curvature, and episodic back-arc rifting in subduction settings: *Geochemistry Geophysics Geosystems*, v. 10, Q05001, doi:10.1029/2008GC002220.
- Wortel, M.J.R., and Spakman, W., 2000, Geophysics—Subduction and slab detachment in the Mediterranean-Carpathian region: *Science*, v. 290, p. 1910–1917, doi:10.1126/science.290.5498.1910.
- Zhong, S.J., Zuber, M.T., Moresi, L., and Gurnis, M., 2000, Role of temperature-dependent viscosity and surface plates in spherical shell models of mantle convection: *Journal of Geophysical Research*, v. 105, p. 11063–11082, doi:10.1029/2000JB900003.

Manuscript received 8 January 2014

Revised manuscript received 17 March 2014

Manuscript accepted 18 March 2014

Printed in USA